2004

Subject Index

AACVPR. *See* American Association of Cardiovascular and Pulmonary Rehabilitation
Abdominal training, electrical stimulation and, 357
Abstracts, AACVPR 2004 Annual Meeting, 342
Access to cardiac rehabilitation, increasing, 171
Active limb mobilization with electrical stimulation, in bed-bound patients with COPD, 283
Acute coronary syndrome, rehabilitated patients after, 38
Acute myocardial infarction, recovery from, scheduled relaxing music sessions and, 343
Adherence to exercise, coronary heart disease and, 348
Aerobic exercise training
  combined with resistance training, in COPD, 359
  effects on aortic wave velocity in obesity, 352
Alternative programs, attendance and graduation patterns, 150
American Association of Cardiovascular and Pulmonary Rehabilitation (AACVPR)
  2004 Annual Meeting, 340
  2004 Annual Meeting abstracts, 342
  Award of Excellence lecture, 146
  consensus statement on outcomes evaluation, 68
  future directions, 146
Anaerobic threshold, effects on chronotropic responses during cardiac pacing, 363
Anxiety, in coronary heart disease, review, 414
Aortic wave velocity, aerobic exercise training and, in obesity, 352
Attendance
  of ethnic minority patients, 308
  and outcomes, 313
  patterns of, 150

Balance assessment, in COPD, 274
Beta-blockers, effects on cardiopulmonary exercise testing with heart failure, 350
Blood glucose, weight management programs and, 349
BODE Index scores, pulmonary rehabilitation programs and, in COPD, 360
Body composition, self-help weight management programs and, 349
Bruce, Robert A., in memoriam, 216
Bypass surgery
  cardiac rehabilitation after
    predicting participation in, 19
    and recovery, 236
    response to, 363
  maximal graded exercise test after, 351
  6-minute walk test after, 351

C-reactive protein, reduction in, 281
CACR. *See* Canadian Association of Cardiac Rehabilitation
Calcium, coronary artery score, combined with Framingham score, 206
California, pulmonary rehabilitation effects on dyspnea, quality of life, and healthcare costs, 52
California Pulmonary Rehabilitation Collaborative Group, 52
California State University, Fullerton Physical Activity Questionnaire, 350
Canada
  metabolic profile and exercise capacity outcomes of cardiac rehabilitation, 405
  referral rate and outcomes of cardiac rehabilitation after cardiac catheterization, 392
Canadian Association of Cardiac Rehabilitation (CACR)
  14th Annual Meeting and Scientific Symposium, 361
  CACR News, 63
  President's Report, 63
Cancer
  exercise during treatment of, 355
  peak exercise response to exercise testing with, 356
Carbon dioxide, VE/VCO_2 slope, prognostic value of, 317
Cardiac catheterization, referral rate and outcomes of cardiac rehabilitation after, 392
Cardiac events, smoking status after, 94
Cardiac pacing, anaerobic threshold effects on chronotropic responses during, 363
Cardiac rehabilitation
  access to, increasing, 171
  after acute coronary syndrome, 38
  adherence to, coronary heart disease and, 348
  alternative approach to delivery of services, 383
  barriers to physical activity maintenance after, 296
  benefits of
    in coronary heart disease, 353
    in men vs women, 45
    older adults, 353
    strain imaging of, 342
  C-reactive protein reduction through, 281
  after CABG surgery, 363
  after cardiac catheterization, referral rate and outcomes of, 392
  with chronic kidney disease, 345
  with combination resistance training and endurance training, outcomes of, 362
  completion rates, 354
  after coronary stenting, 342
  depressive symptoms and, 354
  with diabetes
    benchmarking outcomes for, 348
    gender and risk factor profiles and outcomes in, 361
  dietitian services and patient outcomes in, 352
  employer support and work adjustment in, 362
  ethnic minority patients in, 308
  exercise capacity outcomes, 405
  functional fitness assessment of, in older adults, 34
  gender differences in, 45, 354
  HDL cholesterol response to, 248
  hospitalization for congestive heart failure after, 351
  lipids and lipoproteins after, gender differences in, 257
  maintenance, leisure time physical activity of elderly coronary artery disease patients in, 353

metabolic profile outcomes, 405
for older adults, 245
for older adults, functional fitness assessment of, 34
optimizing risk stratification in, with comorbidity index, 8
outcomes of
evaluation of, 68
exercise training, predicting, 113
gender differences in, 354
outpatient
in managed care organizations, 157
referral to, 165
smoking status after, 94
timeline for peak improvements during, 374
participation in
after bypass surgery, predicting, 19
community education programs and, 352
depressive symptoms and, 354
electronic communication and, 353
perceived exertion in, exercise prescription with, 178
physical function assessment in, 287
and recovery outcomes after CABG, 236
referral for
increasing, 171
review, 171
regional psychological screening program for, 361
sex differences in, 45
social support for, electronic communication and perceived availability of, 353
weight management interventions in, 346
Cardiac rehabilitation programs
alternative, attendance and graduation patterns, 150
inpatient, and chronic respiratory diseases, 359
Internet-based, 343
nearest home, automatic referral to, 171
opportunity knocks, 146
outpatient
hospital-based, outcomes of, 351
patient characteristics, 353
phase 2
patient characteristics, 353
and self-reported health status, 343, 348, 355
physical activity patterns in, 80
Cardiac resynchronization therapy, and exercise capacity, in chronic heart failure, 1
Cardiac risk factor reduction programs, weight reduction in, 348
Cardiopulmonary exercise testing, beta-blockers and, 350
Cardiorespiratory fitness, effects on conduit and resistant artery stiffness in men, 362
Cardiovascular complications, with COPD exacerbation, 356
Cardiovascular disease
exercise prescription with MET-minutes in, 344
obesity and, 197
obstructive sleep apnea syndrome and risk of, 349
WINS VII evaluation among women smokers with, 218
Cardiovascular rehabilitation, predicting risk in, 16
Cardiovascular risk factors, outcomes of cardiac rehabilitation in overweight and obesity, 405
Catheterization, cardiac, referral rate and outcomes of, cardiac rehabilitation after, 392
Center for Medicare and Medicaid Services (CMS),opportunity knocks, 146
CHF. *See* Chronic heart failure; Congestive heart failure
Cholesterol, HDL, response to cardiac rehabilitation, 248
Chronic heart failure
cardiac resynchronization therapy and exercise capacity in, 1
comparison with COPD, 137
Chronic kidney disease, cardiac rehabilitation with, 345
Chronic lung disease, predictors of completion and response to pulmonary rehabilitation in, 360
Chronic obstructive pulmonary disease
combination aerobic and resistance training in, 359
comparison with CHF, 137
endurance training in, 205
exacerbations of, cardiovascular complications with, 356
measures related to severity of, 274
nutritional support for exercise training in, 284
peripheral muscle strength training in, in bed-bound patients, 283
pulmonary rehabilitation programs and BODE Index scores in, 360
self-management intervention for, 282
strength training and functional fitness in, in older patients, 358
supplemental oxygen and exercise training in, 205
UCP3 mRNA and protein content in, 332
unsupported upper limb exercise training in, 422
ventilation-feedback training effects on exercise tolerance in, 358
Chronic respiratory diseases, inpatient cardiac rehabilitation programs and, 359
Chronotropic responses, anaerobic threshold and, 363
Cigarette smoking, and endogenous fibrinolysis in intermittent claudication, 356
Clothing, and workplace physical activity patterns, 350
Community education programs, and cardiac rehabilitation participation, 352
Comorbidity index, optimizing risk stratification with, 8
Comprehensive cardiac risk factor reduction programs, weight reduction in, 348
Comprehensive stress reduction programs, and health scales, 347
Congestive heart failure
death/readmission rate with home-based management, 342
hospitalization for, after cardiac rehabilitation, 351
ventilatory markers during exercise for, 321
Coordination assessment, in COPD, 274
COPD. *See* Chronic obstructive pulmonary disease
Coronary artery bypass graft surgery
cardiac rehabilitation after
predicting participation in, 19
and recovery, 236
response to, 363
maximal graded exercise test after, 351
6-minute walk test after, 351
Coronary artery calcium score, combined with Framingham score, 206
Coronary artery disease
leisure time physical activity of elderly with, 353
resting metabolic rate in, formulas predicting, 346
Coronary events
and metabolic syndrome, in older adults, 353
return to work after, 365
Coronary heart disease
anxiety, PTSD, and depression in, 414
benefits of cardiac rehabilitation in, 353
and cardiac rehabilitation exercise adherence, 348
Coronary revascularization, percutaneous, individualized coronary risk factor intervention in, 259
Coronary risk factor intervention, individualized, 259
Coronary stenting, cardiac rehabilitation after, 342
Costs
future directions, 146
pulmonary rehabilitation effects on, 52
WINS VII evaluation among women smokers with CVD, 218
CVD. *See* Cardiovascular disease
Cycle exercise
leg tourniquets and oxygen uptake during, 128
6-minute cycle test, 344

Death, psychosocial predictors of, 87
Delivery of cardiac rehabilitation services, alternative approach to, 383
Depression
  in coronary heart disease, review, 414
  phase 2 cardiac rehabilitation programs and self-reported health status in, 355
Depressive symptoms, effects in cardiac rehabilitation, 354
Diabetes mellitus
  control of risk factors for vascular disease in, 206
  gender and risk factor profiles and outcomes in, 361
  type 2
    benchmarking outcomes for cardiac rehabilitation with, 348
    response to cardiac rehabilitation after CABG surgery with, 363
    strength training effects on insulin and glucose metabolism in, 282
Diastolic heart failure, 6-minute walk test for women with, 264
Diet
  low-fat vegan, acceptability of, 229
  role in obesity and cardiovascular disease, 197
Dietitian services, and patient outcomes, 352
Disabled older women, resistance training and total energy expenditure in, 344
Drugs, psychotropic, and quality of life, 354
Duke Health Profile (DUKE), comprehensive stress reduction program and, 347
Dysfunctional health scales, comprehensive stress reduction program and, 347
Dyspnea, pulmonary rehabilitation and, 52

Effort, and ventilatory expired gas variables, 317
Electrical stimulation
  active limb mobilization with, in bed-bound patients with COPD, 283
  effects on abdominal strength, endurance, and muscle tone, 357
Electronic communication, effects on perceived availability of social support and participatory intentions, 353
Employer support, in cardiac rehabilitation, 362
Endurance, abdominal, electrical stimulation and, 357
Endurance training
  combined with resistance training, outcomes of, 362
  in COPD, 205
Energy balance markers, self-help weight management programs and, 349
Energy expenditure, total, resistance training and, in disabled older women, 344
Ethnic minority patients, attendance in cardiac rehabilitation, 308
Exercise
  during cancer treatment, 355
  capacity for
    cardiac resynchronization therapy and, in chronic heart failure, 1
    outcomes of cardiac rehabilitation, 405
  cycle, leg tourniquets and oxygen uptake during, 128
  duration of, and postexercise hypotension, 357
  and glucose-insulin response to feeding, 352
  and glycemic response to feeding, 357
  intensity of, and postexercise hypotension, 269
  musical style and performance of, 357
  post-exercise hypotension
    exercise duration and, 357
    exercise intensity and, 269
  prescription for
    comparison of methods, 27
    with MET-minutes, 344
    without graded exercise test, 178
  regular, self-efficacy of physical activity measured during hospitalization and, 362
  resistance
    left ventricular function during, 100
    safety with left ventricular dysfunction, 110
  tolerance to, ventilation-feedback training and, in COPD, 358
Exercise testing
  cardiopulmonary, beta-blockers and, 350
  graded, exercise prescription without, 178
  maximal graded, in CABG patients, 351
  peak response to, in cancer, 356
  6-minute cycle test, 344
  6-minute walk test
    in CABG patients, 351
    for women with diastolic heart failure, 264
Exercise training
  adherence to, coronary heart disease and, 348
  aerobic
    and aortic wave velocity, in obesity, 352
    combined with resistance training, in COPD, 359
  C-reactive protein reduction through, 281
  for congestive heart failure, ventilatory markers during, 321
  effects on oxidative stress, gender differences in, 346
  effects on ventilatory parameters in cardiac patients, 351
  future directions, 146
  nutritional support for, in COPD, 284
  oxygen pulse after, in older adults, 345
  for peripheral arterial disease, pilot intervention, 187
  predicting outcome of, 113
  with self-regulation, 27
  supplemental oxygen in, in COPD, 205
  upper limb, unsupported, in COPD, 422
Exertion, perceived, exercise prescription with, 178

Fasting lipids, weight management programs and, 349
Fat, low-fat vegan diet, acceptability of, 229
Feeding, glycemic responses to
  post-prandial exercise and, 357
  pre-prandial exercise and, 352
Fibrinolysis, endogenous, cigarette smoking and, with intermittent claudication, 356
Framingham score, combined with coronary artery calcium score, 206
Functional effects of COPD, 274
Functional fitness assessment of older cardiac rehabilitation patients, 34
Functional health scales, comprehensive stress reduction program and, 347

Gender differences
  effects on risk factor profiles and outcomes in diabetes mellitus, 361
  in exercise training effects on oxidative stress, 346
  in HDL cholesterol response to cardiac rehabilitation, 248
  in lipids and lipoproteins after cardiac rehabilitation, 257
  in outcomes of cardiac rehabilitation, 354
Glucose blood levels, weight management programs and, 349
Glucose metabolism, strength training and, in type 2 diabetes, 282
Glycemic index, pre-prandial, and glucose-insulin response to feeding, 352
Glycemic responses to feeding
  post-prandial exercise duration and, 357
  pre-prandial exercise and, 352
Graded exercise testing, exercise prescription without, 178
Graduation patterns, 150

Health Care Financing Administration (HCFA), opportunity knocks, 146
Health care providers, perceptions of women's referral to outpatient cardiac rehabilitation, 387
Health maintenance organizations, attendance and graduation patterns, 150
Health-related quality of life
  with ICDs, 324
  pulmonary rehabilitation effects on, 52
Health scales, comprehensive stress reduction program and, 347
Health service utilization, WINS VII evaluation among women smokers with CVD, 218
Health status, self-reported, phase 2 cardiac rehabilitation programs and, 343, 348, 355
Healthcare costs
  future directions, 146
  pulmonary rehabilitation effects on, 52
Heart disease
  community education programs directed to women on, and cardiac rehabilitation participation, 352
  coronary
    benefits of cardiac rehabilitation in, 353
    and cardiac rehabilitation exercise adherence, 348
Heart failure
  beta-blockers and cardiopulmonary exercise testing with, 350
  chronic
    cardiac resynchronization therapy and exercise capacity in, 1
    comparison with COPD, 137
  congestive
    hospitalization for, after cardiac rehabilitation, 351
    ventilatory markers during exercise for, 321
  diastolic, 6-minute walk test for women with, 264
  effort effects on ventilatory expired gas variables in, 317
  phase 1 of oxygen consumption on-kinetics response in, 401
  phase 2 cardiac rehabilitation program effects on self-reported health status after, 343
Heart rate variability, scheduled relaxing music sessions and, 343
High-density lipoprotein cholesterol, response to cardiac rehabilitation, 248
Home-based management, CHF death/readmission rate with, 342
Hospital-based outpatient cardiac rehabilitation programs, outcomes of, 351
Hospitalization
  for congestive heart failure, after cardiac rehabilitation, 351
  physical activity measurement during, self-efficacy of, 362
Hospitals
  CHF death/death admission rate with home-based management, 342
  reduction of utilization of, 282
"Hybrid" model for patient care, 383
Hypotension, postexercise
  exercise duration and, 357
  exercise intensity and, 269

ICDs. *See* Implantable cardioverter defibrillators
Implantable cardioverter defibrillators (ICDs)
  managing patients with, 209
  quality of life with, 324
Individualized coronary risk factor intervention, 259
Insulin
  glucose-insulin response to feeding, effects of pre-prandial exercise and glycemic index on, 352
  strength training and, in type 2 diabetes, 282
Intermittent claudication, cigarette smoking and endogenous fibrinolysis in, 356
Internet-based programs, 343

Journal of Cardiopulmonary Rehabilitation [From the Editor], 67

Kidney disease, chronic, cardiac rehabilitation with, 345

Left ventricular dysfunction, safety of resistance exercise with, 110
Left ventricular function, during strength testing, 100
Leg tourniquets, and oxygen uptake, 128
Leisure time physical activity, of elderly coronary artery disease patients in maintenance cardiac rehabilitation, 353
Lipids
  after cardiac rehabilitation, gender differences in, 257
  fasting, weight management programs and, 349
  HDL cholesterol response to cardiac rehabilitation, 248
Lipoproteins, after cardiac rehabilitation, gender differences in, 257
Literacy, readability of patient education materials and, 121
Literature Update, 205, 281
Low-risk patients, psychosocial predictors of death for, 87
Lower extremity, pilot exercise intervention for, 187
Lung disease, chronic, predictors of completion and response to pulmonary rehabilitation in, 360

Maintenance cardiac rehabilitation, leisure time physical activity of elderly coronary artery disease patients in, 353
Managed care organizations, outpatient cardiac rehabilitation in, 157
Maximal graded exercise tests, in CABG patients, 351
Mechanical ventilation, peripheral muscle strength training in bed-bound patients with COPD on, 283
MEDFICTS questionnaire, 352
Men
  benefits of cardiac rehabilitation for, 45
  cardiorespiratory fitness effects on conduit and resistant artery stiffness in, 362
Messenger RNA (mRNA), UCP-3, in COPD, 332
MET-minutes, exercise prescription with, 344
Metabolic profile, outcomes of cardiac rehabilitation, 405
Metabolic syndrome, coronary events and, in older adults, 353
Minority patients, attendance in cardiac rehabilitation, 308
Minute ventilation (VE), VE/VCO_2 slope, prognostic value of, 317
Mobility assessment, in COPD, 274
Muscle strength, oxygen pulse after exercise training and, 345
Muscle tone, abdominal, electrical stimulation and, 357
Music sessions, scheduled relaxing, and recovery from acute myocardial infarction, 343
Musical style, and exercise performance, 357
Myocardial infarction
  acute, recovery from, scheduled relaxing music sessions and, 343
  psychosocial predictors of death after, 87
Myocardial oxygen demand, scheduled relaxing music sessions and, 343

Nurse-directed home-based management, CHF death/readmission rate with, 342
Nutritional support, for exercise training in COPD, 284

Obesity
  aerobic exercise training effects on aortic wave velocity in, 352
  and cardiovascular disease, 197
  metabolic profile and exercise capacity outcomes of cardiac rehabilitation in, 405
  phase 2 cardiac rehabilitation program effects on self-reported health status in, 348
Obstructive sleep apnea syndrome
  and cardiovascular disease risk, 349
  and physical function, 346
Old-old adults, phase II outpatient cardiac rehabilitation patient characteristics of, 353

Older adults
Cal State Fullerton Physical Activity Questionnaire validity in, 350
cardiac rehabilitation for, 245
benefits of, 353
functional fitness assessment of, 34
with COPD, strength training and functional fitness in, 358
with coronary artery disease, leisure time physical activity of, 353
disabled older women, resistance training and total energy expenditure in, 344
oxygen pulse after exercise training in, 345
recovery after CABG, 236
Optimism, and quality of life with ICDs, 324
Outcomes
AACVPR consensus statement on evaluation of, 68
attendance and, 313
benchmarking, for cardiac rehabilitation with type 2 diabetes, 348
of cardiac rehabilitation after cardiac catheterization, 392
exercise capacity, 405
metabolic profile, 405
timeline for peak improvements during outpatient cardiac rehabilitation, 374
Outpatient cardiac rehabilitation
hospital-based programs, outcomes of, 351
in managed care organizations, 157
phase II programs, patient characteristics of, 353
referral to, 165
smoking status after, 94
timeline for peak improvements during, 374
women's referral to, provider perceptions of, 387
Overweight, metabolic profile and exercise capacity outcomes of cardiac rehabilitation in, 405
Oxidative stress, gender differences in exercise training effects on, 346
Oxygen, supplemental
in exercise training in COPD, 205
pulmonary rehabilitation and, 359
Oxygen consumption (VO_2)
leg tourniquets and, 128
maximal, baseline predictors, 358
peak, prognostic value of, 317
phase 1 on-kinetics response, prognostic value in heart failure, 401
Oxygen demand, scheduled relaxing music sessions and, 343
Oxygen pulse (O2P), after exercise training, in older adults, 345

Pacing, cardiac, anaerobic threshold and chronotropic responses during, 363
Patient education
community education programs, and cardiac rehabilitation participation, 352
readability of materials, 121
Percutaneous coronary revascularization, individualized coronary risk factor intervention in, 259
Performance-based measures of physical function in cardiac rehabilitation, 287
Peripheral arterial disease, pilot exercise intervention for, 187
Peripheral muscle strength training, in bed-bound patients, 283
Phase II cardiac rehabilitation programs
outpatient, patient characteristics of, 353
and self-reported health status, 343, 348, 355
Physical activity
barriers to maintenance after cardiac rehabilitation, 296
Cal State Fullerton Physical Activity Questionnaire, validity in older adults, 350
in cardiac rehabilitation programs, 80
leisure time, of elderly coronary artery disease patients in maintenance cardiac rehabilitation, 353
measurement during hospitalization, self-efficacy of, 362
role in obesity and cardiovascular disease, 197
workplace patterns, clothing and, 350
Physical capacity, and quality of life, 354
Physical function
assessment of, in cardiac rehabilitation, 287
obstructive sleep apnea syndrome and, 346
Positive health expectations, and quality of life with ICDs, 324
Posttraumatic stress disorder, in coronary heart disease, review, 414
Protein, UCP-3, in COPD, 332
Proxy-report of physical function assessment in cardiac rehabilitation, 287
Psychological screening programs, regional, 361
Psychosocial factors, predictors of death, 87
Psychotropic drugs, and quality of life, 354
PTSD. *See* Posttraumatic stress disorder
Pulmonary disease. *See* Chronic obstructive pulmonary disease
Pulmonary rehabilitation
effects on dyspnea, quality of life, and healthcare costs, 52
predictors of completion and response in chronic lung disease, 360
and supplemental oxygen use, 359
Pulmonary rehabilitation programs
and BODE Index scores, in COPD, 360
opportunity knocks, 146

Quality of life
with ICDs, 324
pulmonary rehabilitation and, 52
rehabilitation and, 354

Readability of patient education materials, 121
Referral
automatic, 171
to cardiac rehabilitation after cardiac catheterization, 392
to outpatient cardiac rehabilitation, 165
women's, provider perceptions of, 387
to programs nearest home, 171
of rehabilitated patients after acute coronary syndrome, 38
to rehabilitation, 175
review, 171
Regional psychological screening programs, 361
Rehabilitation. *See also* Cardiac rehabilitation; Pulmonary rehabilitation
and quality of life, 354
residential programs, training responses with self-regulation in, 27
Renal insufficiency, chronic, cardiac rehabilitation with, 345
Residential rehabilitation programs, training responses with self-regulation in, 27
Resistance exercise
left ventricular function during, 100
safety with left ventricular dysfunction, 110
Resistance training
combined with aerobic training, in COPD, 359
combined with endurance training, outcomes of, 362
and total energy expenditure in disabled older women, 344
Respiratory diseases, chronic, inpatient cardiac rehabilitation programs and, 359
Resting heart rate, exercise prescription with, 178
Resting metabolic rate, in coronary artery disease, formulas predicting, 346
Return to work after coronary events, 365
Risk prediction, 16
combination coronary artery calcium score and Framingham score for, 206
Risk stratification, optimizing, with comorbidity index, 8
Rural communities, individualized coronary risk factor intervention in, 259

Screening programs, psychological, regional, 361
Secondary prevention programs, outcomes evaluation of, 68
Self-help weight management programs, effects on body composition and energy balance markers, 349
Self-management intervention, disease-specific, for COPD, 282
Self-paced cycle ergometer testing, 344
Self-regulation, training responses with, 27
Self-report of physical function assessment in cardiac rehabilitation, 287
Self-reported health status, phase 2 cardiac rehabilitation programs and, 343, 348, 355
Service delivery, alternative, 383
Sex differences
  in cardiac rehabilitation, 45
  in HDL cholesterol response to cardiac rehabilitation, 248
6-minute cycle test, 344
6-minute walk test
  in CABG patients, 351
  for women with diastolic heart failure, 264
Skeletal muscle
  endurance training and, in COPD, 205
  strength training effects on insulin signaling in, in type 2 diabetes, 282
  UCP-3 mRNA and protein content, in COPD, 332
Sleep apnea, obstructive
  and cardiovascular disease risk, 349
  and physical function, 346
Slendertone Flex Abdominal Training System, effects of, 357
Small community hospitals, CHF death/death admission rate with home-based management, 342
Smoking, after cardiac events and outpatient cardiac rehabilitation, 94
Smoking cessation interventions
  effectiveness of, 347
  WINS VII evaluation among women smokers with CVD, 218
Social support
  electronic communication and perceived availability of, 353
  and participation in cardiac rehabilitation after bypass surgery, 19
Stenting, coronary, cardiac rehabilitation after, 342
Strain imaging of benefits of cardiac rehabilitation, 342
Strength, abdominal, electrical stimulation and, 357
Strength testing, left ventricular function during, 100
Strength training
  effects on functional fitness in older COPD patients, 358
  effects on insulin and glucose metabolism in type 2 diabetes, 282
  peripheral muscle, in bed-bound patients, 283
Stress
  oxidative, gender differences in exercise training effects on, 346
  posttraumatic, in coronary heart disease, review, 414
Stress reduction programs, and health scales, 347
Supplemental oxygen
  in exercise training in COPD, 205
  pulmonary rehabilitation and, 359
Surgery, bypass
  cardiac rehabilitation after
    predicting participation in, 19
    and recovery, 236
    response to, 363
  maximal graded exercise test after, 351
  6-minute walk test after, 351

Telemedicine, 343
Tobacco cessation interventions
  effectiveness of, 347
  WINS VII evaluation among women smokers with CVD, 218
Total energy expenditure, resistance training and, in disabled older women, 344
Training
  endurance
    combined with resistance training, outcomes of, 362
    in COPD, 205
  exercise
    C-reactive protein reduction through, 281
    nutritional support for, in COPD, 284
    for peripheral arterial disease, pilot intervention, 187
    predicting outcome of, 113
  resistance, combined with endurance training, outcomes of, 362
  with self-regulation, 27
  strength, peripheral muscle, in bed-bound patients, 283
  ventilation-feedback, and exercise tolerance with COPD, 358
Trauma, posttraumatic stress disorder, in coronary heart disease, review, 414

Uncoupling protein-3, mRNA and protein content, in COPD, 332
Upper limb exercise training, unsupported, in COPD, 422

Vascular disease risk factors, in diabetes, 206
Vegan diet, low-fat, acceptability of, 229
Ventilation-feedback training, and exercise tolerance with COPD, 358
Ventilatory expired gas variables, effort effects of, 317
Ventilatory markers, during exercise for CHF, 321
Ventilatory parameters, exercise training and, in cardiac patients, 351

Walking, 6-minute walk test
  in CABG patients, 351
  for women with diastolic heart failure, 264
Weight management
  interventions, effects in cardiac rehabilitation, 346
  programs
    effects on body composition and energy balance markers, 349
    effects on fasting lipid and glucose levels, 349
  reduction, in comprehensive cardiac risk factor reduction programs, 348
Women
  benefits of cardiac rehabilitation for, 45
  community education programs directed to, and cardiac rehabilitation participation, 352
  with diastolic heart failure, 6-minute walk test for, 264
  older, disabled, resistance training and total energy expenditure in, 344
  referral to outpatient cardiac rehabilitation, provider perceptions of, 387
  smokers with CVD, WINS VII evaluation among, 218
Women's Initiative for Nonsmoking (WINS)—VII, 218
Work adjustment, in cardiac rehabilitation, 362
Workplace physical activity patterns, clothing and, 350

Young, phase II outpatient cardiac rehabilitation patient characteristics, 353
Young-old, phase II outpatient cardiac rehabilitation patient characteristics, 353